



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

tion in these plants. Emphasis is laid upon the propagation by tubers, tuberous rootstocks, winter buds, burs, and by fragments of stems. The economic aspect of the genus is inseparable from the ecology of the pond, for it deals principally with the food supply afforded a large number of animals, ranging from the larvae of Diptera to canvasback ducks. The paper is made more valuable by its numerous plates and by an extensive bibliography.—GEO. D. FULLER.

Marine algae in fresh water.—Experimenting with several species of marine algae, Miss BROWN⁴⁰ found that many soon died and disintegrated if placed in fresh water, or even in sea water with a larger admixture of fresh water. Other species, and notably *Enteromorpha intestinalis*, not only endured almost pure fresh water for a period of about 4 weeks, but also seemed to grow more rapidly in waters fresher than those of the sea. The factors involved in this tolerance were not further analyzed.—GEO. D. FULLER.

Thelephoraceae.—In continuing his studies of the Thelephoraceae of North America, BURT⁴¹ presents *Exobasidium*, *Tremellodendron*, *Eichleriella*, and *Sebacina*, with a full historical discussion, synonymy, and citation of stations. The data in reference to the species are as follows: *Exobasidium*, 3 species; *Tremellodendron*, 7 species (2 new species and 3 new combinations); *Eichleriella*, 5 species (2 new species and 3 new combinations); *Sebacina*, 14 species (7 new species and 4 new combinations).—J. M. C.

Species of *Carex*.—MACKENZIE,⁴² in continuing his studies of *Carex*, discusses *C. straminea* and some of its nearest allies, and also describes 8 miscellaneous new species, chiefly western, as follows: *C. festivella*, *C. Egglestonii*, *C. Lunelliana*, *C. bulbostylis*, *C. onustra*, *C. Sheldonii*, *C. exserta*, and *C. rugosperma*.—J. M. C.

Polyporaceae.—OVERHOLTS⁴³ has investigated certain critical forms of Polyporaceae, discussing characters and technique, and presents his conclusions in definitions of the 22 species considered.—J. M. C.

⁴⁰ BROWN, LOLA B., Experiments with marine algae in fresh water. Puget Sound Marine Sta. Publ. 1:31-34. 1915.

⁴¹ BURT, EDWARD ANGUS, The Thelephoraceae of North America. IV and V. Ann. Mo. Bot. Gard. 2:627-658. pl. 21; 731-770. pls. 26, 27. 1915.

⁴² MACKENZIE, K. K., Notes on *Carex*. IX. Bull. Torr. Bot. Club 42:603-621. 1915.

⁴³ OVERHOLTS, L. O., Comparative studies in the Polyporaceae. Ann. Mo. Bot. Gard. 2:667-730. pls. 23-25. 1915.